

**U.S. FISH AND WILDLIFE SERVICE
SPECIES ASSESSMENT AND LISTING PRIORITY ASSIGNMENT FORM**

SCIENTIFIC NAME: *Gallicolumba stairi*

COMMON NAME: Friendly (shy) ground-dove (American Samoa Distinct Population Segment)

LEAD REGION: Region 1

INFORMATION CURRENT AS OF: April 2007

STATUS/ACTION

Species assessment - determined we do not have sufficient information on file to support a proposal to list the species and, therefore, it was not elevated to Candidate status

New candidate

Continuing candidate

Non-petitioned

Petitioned - Date petition received: May 11, 2004

90-day positive - FR date:

12-month warranted but precluded - FR date: May 11, 2005

Did the petition request a reclassification of a listed species?

FOR PETITIONED CANDIDATE SPECIES:

- a. Is listing warranted (if yes, see summary of threats below)? Yes
- b. To date, has publication of a proposal to list been precluded by other higher priority listing actions? Yes
- c. If the answer to a. and b. is "yes," provide an explanation of why the action is precluded.

We find that the immediate issuance of a proposed rule and timely promulgation of a final rule for this species has been, since publication of the last CNOR, and continues to be, precluded by higher priority listing actions (including candidate species with lower LPNs) because most of our national listing budget has been consumed by work on various listing actions to comply with court orders and court-approved settlement agreements, meeting statutory deadlines for petition findings or listing determinations, emergency listing evaluations and determinations, and essential litigation-related, administrative, and program management tasks. We will continue to monitor the status of this species as new information becomes available. This review will determine if a change in status is warranted, including the need to make prompt use of emergency listing procedures. For information on listing actions taken, see the discussion of "Progress on Revising the Lists" in the current CNOR, which can be viewed on our Internet website (<http://www.fws.gov/endangered>).

Y Listing priority change
Former LP: 6
New LP: 9

Date when the species first became a Candidate (as currently defined): November 15, 1994

___ Candidate removal: Former LPN: ___

___ A – Taxon is more abundant or widespread than previously believed or not subject to the degree of threats sufficient to warrant issuance of a proposed listing or continuance of candidate status.

___ U – Taxon not subject to the degree of threats sufficient to warrant issuance of a proposed listing or continuance of candidate status due, in part or totally, to conservation efforts that remove or reduce the threats to the species.

___ F – Range is no longer a U.S. territory.

___ I – Insufficient information exists on biological vulnerability and threats to support listing.

___ M – Taxon mistakenly included in past notice of review.

___ N – Taxon does not meet the Act’s definition of “species.”

___ X – Taxon believed to be extinct.

ANIMAL/PLANT GROUP AND FAMILY: Birds: Family Columbidae (Pigeons and Doves)

HISTORICAL STATES/TERRITORIES/COUNTRIES OF OCCURRENCE: American Samoa (Ofu), Fiji, Tonga, Wallis and Futuna, Independent Samoa

CURRENT STATES/COUNTIES/TERRITORIES/COUNTRIES OF OCCURRENCE: American Samoa (Ofu, Olosega), Fiji, Tonga, Wallis and Futuna, Independent Samoa

LAND OWNERSHIP: All of the land on which the friendly ground-dove occurs in American Samoa is traditionally owned and managed.

LEAD REGION CONTACT: Scott McCarthy, (503) 231-6131, scott_mccarthy@fws.gov

LEAD FIELD OFFICE CONTACT: Pacific Islands Fish and Wildlife Office, Christa Russell, (808) 792-9400, christa_russell@fws.gov

BIOLOGICAL INFORMATION

Species Description

The following description is from Watling (2001). The friendly (shy) ground-dove is an approximately 10.2-inch (26-centimeter) long, medium-sized, brown ground-dove. Males have rufous-brown upperparts with a bronze-green iridescence, the crown and nape are grey and the wings rufous with a purplish lustre. The tail is dark brown. The abdomen and belly are dark

brown olive, while the breast shield is a vinaceous pink with a white border. Females are dimorphic in Fiji and Tonga, but apparently not in Independent Samoa and American Samoa, where only the pale phase occurs. The pale phase is similar to males but rather duller. Immature birds are similar to adults but are uniformly brown.

Taxonomy

The genus *Gallicolumba* is distributed throughout the Pacific and Southeast Asia. The genus is represented in the oceanic Pacific by six species. Three species are endemic to Micronesian islands or archipelagos, two are endemic to island groups in French Polynesia, and *Gallicolumba stairi* is endemic to Samoa, Tonga, and Fiji (Sibley and Monroe 1990, 1993). Some authors recognize two subspecies of the friendly ground-dove: one, slightly smaller, in the Samoan archipelago (*G. s. stairi*), and the other in Tonga and Fiji (*G. s. vitiensis*; Mayr 1945). However, morphological differences between the two are slight (Watling 2001), and no genetic or other studies have validated the existence of separate subspecies. We have carefully reviewed the available taxonomic information (Mayr 1945; Sibley and Monroe 1990, 1993; Watling 2001) and have concluded the species is a valid taxon.

Habitat/Life History

In American Samoa, the friendly ground-dove is typically found on or near steep, forested slopes, particularly those with an open understory and fine scree or exposed soil (U.R. Tulafono, Department of Marine and Wildlife Resources (DMWR), pers. comm. 2006). Elsewhere the species is known to inhabit brushy vegetation and bamboo thickets on offshore islands and forest habitats on large, high islands (Watling 1982; Clunie 1999). The only life history information available for this species is that it typically builds a nest of twigs several feet from the ground, lays one or two white eggs, and is an omnivorous feeder (Watling 1982; Clunie 1999).

Historical and Current Range/Distribution

In American Samoa, the species was first reported on Ofu in 1976 (Amerson *et al.* 1982), and has been recorded infrequently on Ofu and more commonly on Olosega since the mid 1990s (J. Seamon, DMWR, pers. comm. 2004a; U.R. Tulafono, pers. comm. 2006). The American Samoan population of friendly ground-dove currently is distributed in American Samoa on the islands of Ofu and Olosega (Manua Group).

Population Estimates/Status

The friendly ground-dove is uncommon or rare throughout its range (Steadman and Freifeld 1998; Schuster *et al.* 2000; Freifeld *et al.* 2001; Watling 2001), except for on some small islands in Fiji (Watling 2001); however, this species is not known to be monitored except in American Samoa. Engbring and Ramsey (1989) described the population on Ofu as “very small,” but did not attempt a population estimate. More than 10 ground-doves have been caught on Olosega between 2001 and 2004, suggesting that numbers here are greater than on Ofu, but birds may move between the two islands (J. Seamon, pers. comm. 2004a), which once were a single land mass and are today connected by a causeway that is roughly 492 feet (150 meters) long. No current population estimate is available; the secretive habits of this species make monitoring difficult. Monitoring surveys over the last “5 or so” years do not, however, suggest any change in the relative abundance of the friendly ground-dove (J. Seamon, pers. comm. 2004b).

DISTINCT POPULATION SEGMENT (DPS)

The definition of “species” in section 3(15) of the Endangered Species Act (Act) includes any distinct population segment(s) of any species of vertebrate fish or wildlife that interbreed when mature. For a population to be listed under the Act as a distinct vertebrate population segment, three elements are considered: 1) the discreteness of the population segment in relation to the remainder of the species to which it belongs, 2) the significance of the population segment to the species to which it belongs, and 3) the population segment’s conservation status in relation to the Act’s standards for listing (i.e., is the population segment, when treated as if it were a species, endangered or threatened?) (61 FR 4722).

The species *Gallicolumba stairi* is found in Fiji, Tonga, Wallis and Futuna, Independent Samoa, and American Samoa. The status of the species as a whole is not monitored closely throughout its range, but based on available information, the friendly ground-dove persists in very small numbers in Independent Samoa (Schuster *et al.* 2000; Freifeld *et al.* 2001), and is considered to be among the most endangered of native Samoan bird species (Watling 2001). In Tonga, the species occurs primarily on small, uninhabited islands and in one small area of a larger island (Steadman and Freifeld 1998; Watling 2001). In Fiji, *G. stairi* is thought to be widely distributed but uncommon on large islands and relatively common on some small islands (Watling 2001).

The available information indicates that distinct populations of *Gallicolumba stairi*, a cryptic, understory-dwelling dove not noted for long-distance dispersal, are definable. The distinct population segment of friendly ground-dove in American Samoa is discrete in relation to the remainder of the species as a whole. The genus *Gallicolumba* is widely distributed in the Pacific, but populations of individual species are today restricted to a subset of islands (often small, offshore islets) in any archipelago where they occur, or even to limited areas of single islands (e.g., Steadman and Freifeld 1998; Freifeld *et al.* 2001; Watling 2001). Therefore, populations of species such as *G. stairi*, which occurs in three archipelagos, have become isolated from each other. The population segment of this species in American Samoa (on Ofu and Olosega islands), is therefore distinct based on geographic and distributional isolation from the friendly ground-dove populations in Independent Samoa, Tonga, and Fiji. The distance between the U.S. population and the nearest population, in Samoa, is roughly 100 miles (mi) (161 kilometers (km)). The American Samoan island of Tutuila lies between these two populations; no Tutuila records of friendly ground-dove exist.

A population segment is considered “significant” if its loss would constitute a significant gap in the range of the taxon. The American Samoa population of the friendly ground-dove represents the easternmost distribution of this species. The loss of this population would truncate the species’ range by approximately 100 mi (161 km), or approximately 15 percent of the linear extent of its range, which trends southwest-to-northeast from Fiji to American Samoa. Given the rarity with which this species appears in survey data from Independent Samoa, loss of the American Samoa population may decrease the likelihood of the friendly ground-dove’s long-term persistence in the Samoan archipelago. Southwest of Samoa, the next location where this species is known to occur is the Vava`u Group in Tonga, roughly 350 mi (563 km) away. Unlike

other Pacific Island columbids, this species does not fly high above the canopy; it is an understory species that forages largely on the ground and nests near the ground (Watling 2001). Therefore, the likelihood of recolonization over large distances is low.

Based on the discreteness and significance of the American Samoa population of the friendly ground-dove, the U.S. Fish and Wildlife Service (Service) considers this population to be a distinct vertebrate population segment which warrants review for listing under the Act. The distinct population segment of the friendly ground-dove faces severe but non-imminent threats. See SUMMARY OF THREATS and Rationale for Listing Priority Number, below.

THREATS

A. The present or threatened destruction, modification, or curtailment of its habitat or range.

Clearing of lowland rainforests has been implicated as a limiting factor for these populations. At present, none of the land-clearing or development projects proposed for Ofu or Olosega have been approved or initiated in areas known to be frequented by the friendly ground-dove (U.R. Tulafono, pers. comm. 2006).

B. Overutilization for commercial, recreational, scientific, or educational purposes.

Incidental shooting of this species by hunters may be an important threat. But despite its name, the friendly ground-dove is actually shy and secretive, so it is not likely that hunting is a primary threat.

C. Disease or predation.

Nest predation by rats (*Rattus* sp.) is an important threat to many island birds (Atkinson 1977, 1985), especially ground-nesting species (Moors and Atkinson 1984; Bertram and Nagorsen 1995; Flint 1999; Zino *et al.* 2001). For example, black rats (*R. rattus*) were responsible for the near extirpation of the burrow-nesting Galapagos petrel on Floreana Island (Cruz and Cruz 1987), and for the extinction of the ground-nesting Laysan rail (*Porzana palmeri*), which had been translocated to Midway Atoll prior to the loss of the Laysan population (Fisher and Baldwin 1946). Rats thus may play a role in limiting populations of the friendly ground-dove, although no specific data from American Samoa exist with which to test this hypothesis. Feral cats (*Felis domesticus*) have been observed in remote areas known to be frequented by ground-doves and may prey on friendly ground-doves and other ground-nesting species. Recent investigations suggest that avian malaria may be indigenous and non-pathogenic in American Samoa, and therefore is unlikely to limit bird populations there (Jarvi *et al.* 2003; J. Seamon, pers. comm. 2004a). Although other blood parasites are common in many bird species in American Samoa, none have been reported to date in *G. stairi* samples (Atkinson *et al.* 2006). No information is available about the effects of other avian diseases on this species.

D. The inadequacy of existing regulatory mechanisms.

The inadequacy of existing regulatory mechanisms does not appear to be a threat to this species. Hunting bans are ongoing in American Samoa, and the American Samoa Government Department of Marine and Wildlife Resources has statutory authority to enforce these bans as well as authority to restrict development and habitat modification that may affect species such as

the friendly ground-dove. Poaching is not currently considered a threat to the friendly ground-dove (J. Seamon, pers. comms. 2004a, b).

E. Other natural or manmade factors affecting its continued existence.

Small populations are particularly sensitive to thresholds in a range of environmental and demographic parameters (Pimm *et al.* 1988; Mangel and Tier 1994; Meffe and Carroll 1997). This small population may be at risk of extinction because of the low number of individuals and the high frequency of catastrophic events such as hurricanes. Although severe storms are a natural disturbance with which this species has coexisted for millennia, such storms may affect habitat and food resources for birds and thus increase the threats to a population already suffering predation by non-native mammals. For example, Hurricanes Heta (in January 2004) and Olaf (in February 2005) virtually destroyed suitable habitat at one of the areas on Olosega where this species was most frequently encountered; detections of ground-doves in other areas subsequently increased, suggesting they had moved from the area affected by the storms (J. Seamon, pers. comm. 2005; U.R. Tulafono, pers. comm. 2006). Inbreeding and/or reduced likelihood of locating mates are also potential threats for small populations (Thevenon and Couvet 2002; Frankham 2003).

CONSERVATION MEASURES PLANNED OR IMPLEMENTED

Ongoing bans on hunting native species in American Samoa provide protection for this species, and its status is monitored through regular surveys (described below). In addition, DMWR has an ongoing rat-trapping project at 2 sites on Ofu, which likely benefits the friendly ground-dove (J. Seamon, pers. comm. 2004a).

SUMMARY OF THREATS: The American Samoa Distinct Population Segment of the friendly ground-dove is suspected to be threatened by predation by nonnative mammals, and by natural catastrophes such as hurricanes, which pose a threat due to the small population size and limited distribution of this taxon. Based on our evaluation of these ongoing threats, we conclude that there is sufficient information to develop a proposed listing rule for this species. We find that this species is warranted for listing throughout all its range, and, therefore, find that it is unnecessary to analyze whether it is threatened or endangered in a significant portion of its range.

For species that are being removed from candidate status:

___ Is the removal based in whole or in part on one or more individual conservation efforts that you determined met the standards in the Policy for Evaluation of Conservation Efforts When Making Listing Decisions (PECE)?

RECOMMENDED CONSERVATION MEASURES

- Determine population numbers and status of the species
- Conduct studies on basic life history of the species
- Control predators (rats, cats) at known colonies
- Determine other limiting factors and implement control measures

LISTING PRIORITY

THREAT			
Magnitude	Immediacy	Taxonomy	Priority
High	Imminent	Monotypic genus	1
		Species	2
		Subspecies/population	3
	Non-imminent	Monotypic genus	4
		Species	5
		Subspecies/population	6*
Moderate to Low	Imminent	Monotypic genus	7
		Species	8
		Subspecies/population	9
	Non-imminent	Monotypic genus	10
		Species	11
		Subspecies/population	12

Rationale for listing priority number:

Magnitude:

The magnitude of threats facing the American Samoa Distinct Population Segment of the friendly ground-dove is moderate. The threats to this species from predation by nonnative mammals and its low numbers and limited distribution, are ongoing, but do not appear to be of high magnitude, given the fact that the monitoring surveys have not shown any change in the relative abundance of the friendly ground-dove, at least in the last several years (J. Seamon, pers. comm. 2004b).

Imminence:

Threats to the friendly ground-dove are imminent because they are ongoing.

Rationale for Change in Listing Priority Number:

Continued monitoring of this species suggests that it is not in imminent danger of extinction in American Samoa from predation or threats associated with small population size, although these threats are ongoing. Therefore, we have changed the Listing Priority Number for the American Samoa Distinct Population Segment from 6 to 9.

Yes Have you promptly reviewed all of the information received regarding the species for the purpose of determining whether emergency listing is needed?

Is Emergency Listing Warranted? No. The species does not appear to be appropriate for emergency listing at this time because the immediacy of the threats is not so great as to imperil a significant proportion of the taxon within the timeframe of the routine listing process. If it

becomes apparent that the routine listing process is insufficient to prevent significant losses that may result in this species' extinction, then the emergency rule process for this species will be initiated. We will continue to monitor the status of the friendly ground-dove as new information becomes available. This review will determine if a change in status is warranted, including the need to make prompt use of emergency listing procedures.

DESCRIPTION OF MONITORING

The Department of Marine and Wildlife Resources (DMWR), an agency of the American Samoa Government, is responsible for monitoring the friendly ground-dove and is the sole source of current, on-the-ground information about wildlife in the Territory. This agency is funded through the U.S. Fish and Wildlife Service's Federal Assistance program for wildlife restoration on an annual basis to monitor and manage the fish and wildlife resources of the Territory of American Samoa, and the Service requests annual updates from DMWR on the status of candidate species. We also reviewed current scientific literature to seek new published information about the species in Samoa and elsewhere in its range, and we sent our most current information to regional and species experts for review. We received the latest update from DMWR on August 31, 2006; other queries and literature review yielded no new information. The friendly ground-dove is a secretive species and somewhat difficult to monitor. Several survey methods are therefore employed at a wide array of sites as part of regular surveys of birds in the Manua Islands. Surveys on Ofu and Olosega include annual visual surveys at six sites, biannual auditory surveys at four sites, and mist-net (mark-recapture) surveys at five sites at least annually. These surveys, which have been conducted for the past several years, have so far not yielded evidence of significant change in the relative abundance of the friendly ground-dove (J. Seamon, pers. comm. 2004b). The rate of detection of this secretive bird is insufficient to develop an estimate of total population size. However, the methods, frequency, and geographic coverage of monitoring surveys for this species are adequate to detect changes in distribution and trends in relative abundance over time.

This species is classified as vulnerable in the International Union for Conservation of Nature and Natural Resources (IUCN) Red Data List database (IUCN 2006), and is included in the Comprehensive Strategy for Wildlife Conservation in American Samoa (DMWR 2006).

COORDINATION WITH STATES

The latest assessment form was sent to Ray Tulafono, the Director of the American Samoa Department of Marine and Wildlife Resources on July 13, 2006. He provided additional information and comments on August 31, 2006, and these were incorporated into the form.

LITERATURE CITED

- Amerson, A.B., Jr., W.A. Whistler, and T.D. Schwaner. 1982. Wildlife and wildlife habitat of American Samoa. II. Accounts of flora and Fauna. U.S. Fish and Wildlife Service. 151 pp.
- Atkinson, C.T., R.C. Uzzurum, J.O. Seamon, A.F. Savage, and D.A. LaPointe. 2006. Hematozoa of forest birds in American Samoa – evidence for a diverse, indigenous

- parasite fauna from the South Pacific. *Pacific Conservation Biology*, Vol. 12:229-237.
- Atkinson, I.A.E. 1977. A reassessment of factors, particularly Rattus rattus L., that influenced the decline of endemic forest birds in the Hawaiian Islands. *Pacific Science* 31:109-133.
- Atkinson, I.A.E. 1985. The spread of commensal species of Rattus to oceanic islands and their effects on island avifaunas. Pages 35-81 in P. J. Moors, ed., *Conservation of island birds*. Tech. Publ. No. 3, Int. Coun. Bird Preserv., Cambridge, England.
- Bertram, D. and D. Nagorsen. 1995. Introduced rats Rattus spp. on the Queen Charlotte Islands: implications for seabird conservation. *Canadian Naturalist* 10: 6-10.
- Clunie, F. 1999. *Birds of the Fiji bush*. Fiji Museum, Suva, Fiji. 142 pp.
- Cruz, F. and J.B. Cruz. 1987. Control of black rats (Rattus rattus) and its effect on nesting dark-rumped petrels in the Galapagos Islands. *Vida Silvestre Neotropical* 1: 3-13.
- Department of Marine and Wildlife Resources. 2006. A comprehensive strategy for wildlife conservation in American Samoa (R.C.B. Utzurrum, J.O. Seamon, and K. Schletz Sali, authors). DMWR, Pago Pago. 109 pp.
- Engbring, J., and F.L. Ramsey. 1989. A 1986 survey of the forest birds of American Samoa. U.S. Fish and Wildlife Service. 145 pp.
- Fisher, H.I. and P.H. Baldwin. 1946. War and the birds of Midway Atoll. *Condor* 48: 3-15.
- Flint, E. 1999. Status of seabird populations and conservation in the tropical island Pacific. In L.P. Eldredge, P. Holtus, and J. Maragos (eds.), *Marine and coastal biodiversity in the tropical island Pacific region: population, development, and conservation priorities*, vol. 2. East-West Center, Honolulu, Hawaii. 32 pp. + Appendices.
- Frankham, R. 2003. Genetics and conservation biology. *C.R. Biologies*, 326 Suppl 1: S22-9.
- Freifeld, H.B., D.W. Steadman, and J.K. Sailer. 2001. Observations of birds on offshore islands in Samoa. *Journal of Field Ornithology* 72:72-85.
- IUCN 2006. 2006 IUCN Red List of Threatened Species. www.iucnredlist.org. Downloaded on 15 February 2007.
- Jarvi, S.I., M.E.M. Farias, H. Baker, H.B. Freifeld, P.E. Baker, E. VanGelder, J.G. Massey, C. T. Atkinson. 2003. Detection of avian malaria (Plasmodium sp.) in native land birds of American Samoa. *Conservation Genetics* 4: 629-637.
- Mangel, M. and C. Tier. 1994. Four facts every conservation biologist should know about persistence. *Ecology* 75: 607-614.

- Mayr, E. 1945. Birds of the Southwest Pacific. The Macmillan Company, New York, NY. 316 pp.
- Meffe, G.K., and C.R. Carroll. 1997. Principles of Conservation Biology, 2nd Ed. Sinauer Associates, Sunderland, MA. 729 pp.
- Moors, P.J. and I.A.E. Atkinson. 1984. Predation on seabirds by introduced animals, and factors affecting its severity. Pp. 667-690 in J.P. Croxall, P.G.H. Evans, and R.W. Schreiber (eds.), Status and conservation of the world's seabirds, Int. Coun. Bird Preserv. Tech. Bull. No. 2, Cambridge, U.K.
- Pimm, S., H. L. Jones, and J. Diamond. 1988. On the risk of extinction. American Naturalist 132(6): 757-785.
- Schuster, C., A. Whistler, and T.S. Tuaillemafuna. 2000. The conservation of biological diversity in upland ecosystems of Samoa. Division of Environment and Conservation, Apia, Samoa. 164 pp.
- Sibley, G. and B.L. Monroe, Jr. 1990. Distribution and Taxonomy of Birds of the World. Charles Yale University Press, New Haven and London. 1111 pp.
- Sibley, G. and B.L. Monroe, Jr. 1993. Supplement to Distribution and Taxonomy of Birds of the World. Yale University Press, New Haven and London. 108 pp.
- Steadman, D.W. and H. B. Freifeld. 1998. Distribution and relative abundance of landbirds in the Vava'u group, Kingdom of Tonga. Condor 100: 609-628.
- Thévenon, S. and D. Couvet. 2002. The impact of inbreeding depression on population survival depending on demographic parameters. Animal Conservation 5: 53-60
- Watling, D. 1982. Birds of Fiji, Tonga, and Samoa. Millwood Press, Wellington, New Zealand. 176 pp.
- Watling, D. 2001. Birds of Fiji and Western Polynesia. South Pacific Regional Environmental Programme, Apia, Samoa. 272 pp.
- Zino, F., P. Oliviera, S. King, A. Buckle, M. Biscoito, H. Costa Neves, A. Vasconcelos. 2001. Conservation of Zino's petrel Pterodroma madeira in the archipelago of Madeira. Oryx 35: 128-135.

Personal Communications and *in litt.*:

- Seamon, J.O. Department of Marine and Wildlife Resources. 2004a. Electronic mail message

to Holly Freifeld, U.S. Fish and Wildlife Service, regarding candidate status updates for American Samoa, dated May 31, 2004.

Seamon, J.O. Department of Marine and Wildlife Resources. 2004b. Electronic mail message to Holly Freifeld, U.S. Fish and Wildlife Service, regarding likelihood of poaching and population estimates for the friendly ground-dove, dated June 4, 2004.

Seamon, J.O. Department of Marine and Wildlife Resources. 2005. Electronic mail message to Holly Freifeld, U.S. Fish and Wildlife Service, regarding re-request for assistance: Am. Samoa birds candidate assessments, dated September 20, 2005.

Tulafono, U.R. Department of Marine and Wildlife Resources. 2006. Letter to Patrick Leonard, U.S. Fish and Wildlife Service in response to request for review of the draft candidate assessment forms, dated August 31, 2006.

APPROVAL/CONCURRENCE: Lead Regions must obtain written concurrence from all other Regions within the range of the species before recommending changes, including elevations or removals from candidate status and listing priority changes; the Regional Director must approve all such recommendations. The Director must concur on all resubmitted 12-month petition findings, additions or removal of species from candidate status, and listing priority changes.

Approve: _____
Regional Director, Fish and Wildlife Service Date

Concur: _____
Director, Fish and Wildlife Service Date

Do not concur: _____
Director, Fish and Wildlife Service Date

Director's Remarks:

Date of annual review: _____ Date: April 6, 2007
Conducted by: Karen Rosa, Pacific Islands FWO
Biologist, Prelisting and Listing Program

Comments:
PIFWO Review

Reviewed by: Christa Russell Date: April 13, 2007
Prelisting and Listing Program Coordinator

Gina Shultz Date: April 13, 2007
Assistant Field Supervisor,
Endangered Species

Patrick Leonard Date: April 13, 2007
Field Supervisor

(Revised 8/12/05)